arXiv:1401.6656v7 [math.NT] 22 Sep 2014

GAUSSIAN MERSENNE PRIMES OF THE FORM $x^2 + dy^2$

SUSHMA PALIMAR AND AMBEDKAR DUKKIPATI

Abstract. In this paper we study Gaussian ring $\mathbb{Z}[i]$ with a focus on representing Gaussian Mersenne primes G_p in the form $x^2 + 7y^2$. Interestingly when such a form exists, one can observe that, $x \equiv \pm 1 \pmod 8$ and $y \equiv 0 \pmod 8$. To prove this property of Gaussian Mersenne primes, we show that Gaussian Mersenne primes splits completely in the cyclic quartic unramified extension of $\mathbb{Q}(\sqrt{-14})$ and have a trivial Artin symbol in this extension. We generalize this result for $d \equiv 7 \pmod{24}$. We also attempt to give an alternate proof using Artin's reciprocity law, which was earlier given by H. W. Lenstra and P. Stevenhagen to prove a similar property on ordinary Mersenne Primes.

[1]

1. Introduction

The study of prime numbers in the form $x^2 + dy^2$ for a fixed integer $d > 0$ is ancient and formal study of this problem began with Fermat, later continued by Euler, which led him to discover quadratic reciprocity and many conjecture on $x^2 + dy^2$ for $d > 3$. The book [6] is a good reference for this. Study of Mersenne primes in the form $x^2 + dy^2$, for $d \in [2, 48]$ is carried out in detail in [12]. In [10] Lenstra and Stevenhagen prove an interesting property of Mersenne primes which was first observed by Lemmermeyer. This observation is given in the form of Theorem below. Here authors make use of Artin's reciprocity to demonstrate this property of Mersenne primes.

Theorem 1.1. *Let M_p be a Mersenne prime with $p \equiv 1 \pmod 3$. Write $M_p = x^2 + 7y^2$ with $x, y \in \mathbb{Z}$. Then x is divisible by 8.*

In [14], authors call $\alpha = (2 + \sqrt{2})$ and the norm $N(\frac{\alpha^p - 1}{\alpha - 1})$ is found. In this case whenever the norm $N(\frac{\alpha^p - 1}{\alpha - 1})$ is a rational prime, then those primes are quadratic residues of 7 hence can be written in the form $x^2 + 7y^2$ and it is proved using Artin's reciprocity that, x is divisible by 8 and $y \equiv \pm 3 \pmod 8$.

In Section 2 we study Gaussian Mersenne primes in the form $x^2 + dy^2$ with a special focus on the form $x^2 + 7y^2$. Here we prove that whenever $G_p = x^2 + 7y^2$ then $x \equiv \pm 1 \pmod 8$ and $y \equiv 0 \pmod 8$ and generalize this result for all $d \equiv 7 \pmod{24}$ in the next section.

[1]Research by the first author is supported by National Board for Higher Mathematics-Post Doctoral Fellowship, Government of India.

1.1. **Gaussian Mersenne Primes.** Gaussian Mersenne primes and their primality test was first studied by Berrizbeitia and Iskra in [1]. A Gaussian Mersenne number is an element of $\mathbb{Z}[i]$ given by $\mu_p = (1 \pm i)^p - 1$, for some rational prime p. The Gaussian Mersenne norm $G_p = 2^p - \left(\frac{2}{p}\right) 2^{\frac{p+1}{2}} + 1$ is the norm of μ_p, $N(\mu_p)$. If G_p is a rational prime then we call G_p a Gaussian Mersenne prime.

1.2. **Some results from Class Field Theory.** We take a short and quick revision of class field theory to fix notations. Gal(L/K) is the Galois group of the field extension $K \subset L$. $\mathcal{O}_K$, the ring of algebraic integers in a finite extension K of $\mathbb{Q}$. We denote $I_K(\Delta(L/K))$ to be the group of all fractional ideals of $\mathcal{O}_K$ prime to $\Delta(L/K)$. The *Artin symbol* is denoted by $((L/K)/\mathfrak{P})$, for $\mathfrak{P}$, a prime of L containing a prime $\mathfrak{p}$ of $\mathcal{O}_K$. Then for any α in $\mathcal{O}_L$

$$\left(\frac{L/K}{\mathfrak{P}}\right)(\alpha) \equiv \alpha^{N(\mathfrak{p})} \pmod{\mathfrak{P}},$$

When $K \subset L$ is an Abelean extension, the Artin symbol can be written as $((L/K)/\mathfrak{p})$.

Definition 1.2. *Let K be a number field and $\mathfrak{m}$ a modulus of K. We define $P_{K,1}(m)$ to be the subgroup of $I_K(m)$ generated by the principal ideals $\alpha.\mathbb{Z}_K$, with $\alpha \in \mathbb{Z}_K$, satisfying $\alpha \equiv 1 \bmod \mathfrak{m}_0$, where $\mathfrak{m}_0$ is a nonzero $\mathcal{O}_K$ ideal and $\sigma(\alpha) > 0$ for all infinite primes σ of K dividing $\mathfrak{m}_\infty$, is a product of distinct real infinite primes of K. Thus Cl_m is the group $I_K(m)/P_{K,1}(m)$.*

Given an order $\mathcal{O}$ in a quadratic field K, the index $f = [\mathcal{O} : \mathcal{O}_K]$ is called conductor of the order. The discriminant of $\mathcal{O}$ is $D = f^2 d_k$, d_k is the discriminant of the maximal order $\mathcal{O}_K$.

Definition 1.3. *Let K be a number field and $R = R_m(K)$ the ray class field of K with modulus m. The ray class field of conductor $\mathfrak{m} = (1)$ is the Hilbert class field $H = H_1$ of K.*

Definition 1.4 (Ring Class Field)**.** *If $\mathcal{O}$ is the order of $\mathcal{O}_K$ of conductor f, we define ring class group to be $I_K(f\mathcal{O}_K)/P_{K,\mathbb{Z}}(f\mathcal{O}_K)$, which is naturally isomorphic to the group of ideals prime to f modulo the principal ideals prime to f. The ring class field of the maximal order $\mathcal{O}_K$ is the Hilbert class field of K.*

Here, the bottom group $P_{K,\mathbb{Z}}(f\mathcal{O}_K)$ consists of the principal ideals which admit a generator α congruent to some integer a, relatively prime to f, i.e.,

$$\alpha \equiv a \pmod{f\mathcal{O}_K}$$

We have $m = [\mathcal{O}_K : \mathcal{O}] = 1, 2, 3, 4, 6$ if and only if $P_{K,1}(m) = P_{K,\mathbb{Z}}(m)$. This is because, $m = 1, 2, 3, 4, 6$ is nothing but $\phi(m) \leq 2$. And these are the cases where the global units 1 and –1 'fill up' $(\mathbb{Z}/m\mathbb{Z})^*$, and the ray class field equals the ring class field.

2. Gaussian Mersenne primes as $x^2 + dy^2$

In this section we derive two properties of Gaussian Mersenne primes and prove our main result that, whenever $G_p = x^2 + 7y^2$ then $x \equiv \pm 1 \pmod 8$ and $y \equiv 0 \pmod 8$.

Definition 2.1. *We define F_d to be the quadratic form $x^2 + dy^2 \in \mathbb{Z}[x, y]$. Let p be a prime then p is represented by F_d if there exists $x > 0$ and $y > 0$ in $\mathbb{Z}$ such that $p = x^2 + dy^2$.*

Proposition 2.2. *Let $d \equiv 3 \bmod 4$ be a square-free integer. Suppose $L = R_2(\mathbb{Q}(\sqrt{-d}))(\sqrt{2}) = H(\mathbb{Q}(\sqrt{-2d}))$. Let G_p be a Gaussian Mersenne prime unramified in L then F_d represents G_p if and only if F_{2d} represents G_p.*

This property of Gaussian Mersenne primes stated in Proposition 2.2 is obeyed by Mersenne primes too, and the proof is similar to the proof given in [12].

Proof. We have $G_p = 2^p - \left(\frac{2}{p}\right) 2^{\frac{p+1}{2}} + 1$. For $p > 3$, $G_p \equiv 1 \bmod 8$. Hence, G_p splits completely in $\mathbb{Q}(\sqrt{2})$ and the prime G_p is unramified in L, which implies $G_p \nmid 2d$. We have, $[\mathcal{O}_K : \mathbb{Z}[\sqrt{-d}]] = 2$, since $d \equiv 3 \pmod 4$ is a positive square-free integer. Now, by using the fact G_p splits in $\mathbb{Q}(\sqrt{2})$ we get F_d represents G_p if and only if G_p splits completely in $R_2(\mathbb{Q}(\sqrt{-d}))(\sqrt{2})$. For $K = \mathbb{Q}(\sqrt{-2d})$ we have $[\mathcal{O}_K : \mathbb{Z}[\sqrt{-2d}] = 1$. Thus, F_{2d} represents G_p if and only if G_p splits completely in $H(\mathbb{Q}(\sqrt{-2d}))$. By assumption we have $R_2(\mathbb{Q}(\sqrt{-d}))(\sqrt{2}) = H(\mathbb{Q}(\sqrt{-2d}))$ hence the result. □

The property of Gaussian Mersenne primes in Proposition 2.3 is unique only to Gaussian Mersenne primes.

Proposition 2.3. *Let $d \equiv 1 \bmod 4$ be a square-free integer. Suppose $L = H(\mathbb{Q}(\sqrt{-d})) = H(\mathbb{Q}(\sqrt{-2d}))$. Let G_p be a Gaussian Mersenne prime unramified in L then F_d represents G_p if and only if F_{2d} represents G_p, which is not the case for usual Mersenne primes.*

Proof. Proof follows from the previous proposition, except for the fact, $[\mathcal{O}_K : \mathbb{Z}[\sqrt{-d}]] = 2$, which is 1 in the present case, as $d \equiv 1 \bmod 4$. □

2.1. Gaussian Mersenne primes as $x^2 + 7y^2$.

Here we give first few examples of Gaussian Mersenne primes and their representation as $x^2 + 7y^2$.

$$G_7 = 113 = 1 + 7 \cdot 4^2$$

$$G_{47} = 140737471578113 = 5732351^2 + 7 \cdot 3925696^2$$

$$G_{73} = 9444732965601851473921 = 96890022433^2 + 7 \cdot 2854983576^2$$

$$G_{113} = 10384593717069655112945804582584321 =$$

$$792885099381473612 + 7 \cdot 24195412519312600^2$$

One can check that, Gaussian Mersenne prime G_p is $1 \pmod 8$ for all $p > 3$. Also ,

$$\text{if } p \equiv 1 \pmod 6 \text{ then } Gp \equiv 1 \pmod 7$$

and

$$\text{if } p \equiv 5 \pmod 6 \text{ then } Gp \equiv 4 \pmod 7 \text{ for all } p.$$

For $p > 7$, a simple computation shows that, in each case $x \equiv \pm 1 \bmod 8$ and $y \equiv 0 \bmod 8$. Also, from the above table it is not difficult to see that y is exactly divisible by 4. These two observations are proved as a lemma below.

Lemma 2.4. *If G_p is represented in the form x^2+7y^2 then $x \equiv \pm 1 \pmod 8$ and $4|y$.*

Proof. From the above discussion it is clear that, G_p can be written in the form $x^2 + 7y^2$ if and only if $p \equiv \pm 1 \pmod 8$. Also, $G_p = 2^p - 2^{\frac{p+1}{2}} + 1 \equiv 1 \pmod 8$ for all $p > 3$. We first show that, y is divisible by 4. For this we show that, x is an odd integer and y is an even integer. For if x is even and y is odd, then $x^2 \equiv 0$ or $4 \pmod 8$ and $y^2 \equiv 1 \pmod 8$, which implies $x^2 + 7y^2 \equiv 7$ or $3 \pmod 8$, a contradiction. Hence x should be an odd integer and y should be an even integer. Now, clearly $x^2 \equiv 1 \pmod 8$ and $y^2 \equiv 0$ or $4 \pmod 8$. If $y^2 \equiv 4 \pmod 8$ then $x^2 + 7y^2 \equiv 5 \pmod 8$, again a contradiction. Hence $y \equiv 0 \pmod 4$ and we have $4|y$. Now we show that $x \equiv \pm 1 \pmod 8$. Clearly $x^2 + 7y^2 \equiv x^2 \pmod{16}$. Since x is an odd integer either $x^2 \equiv 1 \pmod{16}$ or $x^2 \equiv 9 \pmod{16}$. If $x^2 \equiv 9 \pmod{16}$ then $G_p \equiv 9 \pmod{16}$, a contradiction as $G_p \equiv 1 \pmod{16}$, for all $p \equiv \pm 1 \pmod 8$. Thus, $x^2 \equiv 1 \pmod{16}$. Hence finally $x \equiv \pm 1 \pmod 8$ and $y \equiv 0 \pmod 4$. □

Now it remains to prove that y is divisible by 8 in the expression $G_p = x^2 + 7y^2$. As a first step we transform the base field $\mathbb{Q}(\sqrt{-7})$ into $\mathbb{Q}(\sqrt{2})$. Then show that Gp splits completely in the cyclic quartic unramified extension of $\mathbb{Q}(\sqrt{-14})$. This is because, $G_p \equiv 1 \pmod 8$ for all $p > 3$ which shows that, G_p is a prime in $\mathbb{Q}$ which splits in $\mathbb{Q}(\sqrt{2})$.

Proposition 2.5. *Let $p > 7$ and $G_p = 2^p - (\frac{2}{p})2^{\frac{p+1}{2}} + 1$ be a Gaussian Mersenne prime. Then for $p \equiv \pm 1 \pmod 8$ the form $G_p = x^2 + 7y^2$ exists. Suppose that there exists a cyclic extension H_4 of $S = \mathbb{Q}(\sqrt{-14})$ with $[H_4 : S] = 4$, $H_4 \subset H(S)$ and $\sqrt{2} \in H_4$. Then G_p splits completely in H_4.*

Proof. Let $K = \mathbb{Q}(\sqrt{-7})$ and let $J = \mathbb{Q}(\sqrt{2})$. As $\sqrt{2} \in H_4$, consider an extension $J = \mathbb{Q}(\sqrt{2}) \subset H_4$ and $Gal(H_4/\mathbb{Q})$ is isomorphic to the dihedral group with 8 elements. Thus, there are two conjugate field extensions of J, say J_1 and J_2 contained in H_4, namely $J_1 = J(\sqrt{-1+2\sqrt{2}})$ and $J_2 = J(\sqrt{-1-2\sqrt{2}})$. Consider the following field diagram.



The discriminant of JK over J is -7. The discriminants $\Delta(J_1/J)$ and $\Delta(J_2/J)$ must be relatively prime. If not there is a prime of J which is ramified in J_1 and J_2. This prime would be ramified in JK, because $\Delta(H_4/JK) = 1$. But then the inertia field of this prime ideals equals J, which cannot be the case since $\Delta(H_4/JK) = 1$, using the result on discriminant of tower of fields, and $-7 = \Delta(J_1/J)\Delta(J_2/J)$. Since G_p splits in $\mathbb{Q}(\sqrt{2})$, write $G_p = v_p \cdot \bar{v_p}$, identifying $v_p = x_1 + \sqrt{2}y_1$ and $\bar{v_p} = x_1 - \sqrt{2}y_1 \in \mathbb{Z}_J$ for some $x_1, y_1 \in \mathbb{Z}$. Clearly $\sigma(v_p) > 0$ and $\sigma(\bar{v_p}) > 0$ for all p and for all σ in $Gal(J/\mathbb{Q})$.

$$\text{Claim: } v_p \equiv \bar{v}_p \equiv 1 \pmod{\Delta(J_i/J)} \text{ for } i = 1, 2 \text{ and for all } p$$

Here we consider two cases $p \equiv 1 \pmod 6$ and $p \equiv 5 \pmod 6$. For $p \equiv 1 \pmod 6$, we have $G_p \equiv 1 \pmod 7$ hence, $v_p \equiv \bar{v}_p \equiv 1 \pmod{\Delta(J_1/J)}$ and $v_p \equiv \bar{v}_p \equiv 1 \pmod{\Delta(J_2/J)}$. Since $\Delta(J_1/J)$ and $\Delta(J_2/J)$ are coprime, we get $v_p \equiv \bar{v_p} \equiv 1 \pmod{\Delta(J_i/J)}$ for $i \in 1, 2$. Applying Artin map,

$$\left(\frac{J_i/J}{v_p}\right) = \left(\frac{J_i/J}{\bar{v_p}}\right) = 1 \text{ for } i \in 1, 2,$$

since v_p and $\bar{v}_p$ are in the kernel. The Galois group of H_4/J is isomorphic to $Gal(J_1/J) \times Gal(J_2/J)$. So

$$\left(\frac{H_4/J}{v_p}\right) = \left(\frac{H_4/J}{\bar{v_p}}\right) = 1.$$

So, G_p splits completely in H_4 for $p \equiv 1 \pmod 6$. For $p \equiv 5 \pmod 6$ to show that Artin symbol corresponding to J_1/J and J_2/J is trivial, the strategy has to be some what different. We know that, for $p \equiv 5 \pmod 6$, $v_p \cdot \bar{v}_p \equiv 1 \pmod 8$, which is a prime in $\mathbb{Q}$ splits in $J = \mathbb{Q}(\sqrt{2})$, and therefore J completed at v_p is nothing but $\mathbb{Q}$ completed at $G_p = v_p \cdot \bar{v}_p$. Since $G_p \equiv 1 \pmod 8$, we have,

$$\left(\frac{-1}{G_p}\right) = 1$$

Now it remains to compute the Legendre symbol $\left(\frac{-7}{G_p}\right)$. If $p \equiv 5 \pmod 6$ then $G_p \equiv 4 \pmod 7$, hence $\left(\frac{G_p}{7}\right) = 1$. Therefore by quadratic reciprocity we have,

$$\left(\frac{-7}{G_p}\right) = 1,$$

and the claim follows. □

From Proposition 2.5 and Lemma 2.4 it is clear that G_p splits completely in the cyclic extension of $\mathbb{Q}(\sqrt{-14})$. Now we are ready to prove our main result that, for $p > 7$ and $p \equiv \pm 1 \pmod 8$ in the representation of G_p as $G_p = x^2 + 7y^2$ for some $x, y \in \mathbb{Z}$, y is divisible by 8.

Theorem 2.6. *Let $p > 7$ and $G_p = 2^p - (\frac{2}{p})2^{\frac{p+1}{2}} + 1$ be a Gaussian Mersenne prime. Then for $p \equiv \pm 1 \pmod 8$ the form $G_p = x^2 + 7y^2$ exists for some $x, y \in \mathbb{Z}$ and y is divisible by 8.*

Proof. Here we make use of the fact that, G_p splits completely in the cyclic extension of $\mathbb{Q}(\sqrt{-14})$, so we refer to Proposition 2.5 for rest of the proof. Consider the following lattice of fields.



It is clear that the Galois group of $H_4/\mathbb{Q}$ is the dihedral group with 8 elements. Here $K = \mathbb{Q}(\sqrt{-7})$, $J = \mathbb{Q}(\sqrt{2})$, $S = \mathbb{Q}(\sqrt{-14})$ and $\omega = \frac{1+\sqrt{-7}}{2}$. Let $\bar{\omega}$ be the conjugate of ω then $\omega\bar{\omega} = 2$ and 2 splits completely in K as $2 = (2, \omega)(2, \bar{\omega})$. Now, K_1 and K_2 are the two conjugate field extensions of K, and J_1 and J_2 are the two conjugate field extensions of J respectively. Discriminant of the extension fields are, $\Delta(S/\mathbb{Q}) = -56$, $\Delta(K/\mathbb{Q}) = -7$, $\Delta(J/\mathbb{Q}) = 8$, $\Delta(JK/S) = 1$, $\Delta(H_4/JK) = 1$, $\Delta(JK/J) = -7$ and $\Delta(J_1/J)\Delta(J_2/J) = -7$. Clearly $\Delta(K_1/K)$ and $\Delta(K_2/K)$ are coprime, and $\Delta(K_1/K) \cdot \Delta(K_2/K) = 8$. Since the discriminants are coprime and K_1 and K_2 are conjugates, $\Delta(K_1/K) = (2, \omega)^3$ and $\Delta(K_2/K) = (2, \bar{\omega})^3$. Now, $\Delta(H_4/\mathbb{Q}) = 8^2$ and $\Delta(K_i/K)|8^2$. Let $\Delta = \Delta(K_1/K)$. Then there is a surjective group homomorphism:

$$Cl_\Delta(K) \to Gal(K_2/K).$$

The group $I_K(\Delta)/P_{K,1}(\Delta) = P_K(\Delta)/P_{K,1}(\Delta)$ is a subgroup of $Cl_\Delta(K)$ and contains the principal prime ideal $\pi = (x + y\sqrt{-7})$ of K. Any prime of $\mathbb{Q}$, which is inert in both J and K splits in S. For example, since $7 \equiv 1 \pmod 3$, the prime 3 of $\mathbb{Q}$ is inert in J and K and splits in S. Thus the decomposition group of a prime $\mathfrak{p}_3$ in H_4 above 3 is S. The prime $3\mathbb{Z}_K$ of K does not split in K_1. Hence, $P_K(\Delta)/P_{K,1}(\Delta)$ maps surjective on $Gal(K_1/K)$. The map from $(\mathbb{Z}_K/\Delta)^*$ to $P_K(\Delta)/P_{K,1}(\Delta)$ is surjective and the group $(\mathbb{Z}_K/\Delta)^*$ is isomorphic to the group $(\mathbb{Z}/8\mathbb{Z})^*$. We have the following group homomorphism,

$$(\mathbb{Z}/8\mathbb{Z})^* \simeq (\mathbb{Z}_K/\Delta)^* \to P_K(\Delta)/P_{K,1}(\Delta) \to Gal(K_1/K).$$

Now $\{\pm 1\}$ is contained in the kernel of the $(\mathbb{Z}_K/\Delta)^* \to P_K(\Delta)/P_{K,1}(\Delta)$ and this map is surjective hence, the kernel of $(\mathbb{Z}_K/\Delta)^* \to P_K(\Delta)/P_{K,1}(\Delta)$

is $\{\pm 1\}$. Thus $P_K(\Delta)/P_{K,1}(\Delta)$ has two elements. So,

$$(\mathbb{Z}/8\mathbb{Z})^* \simeq (\mathbb{Z}_K/\Delta)^* \to P_K(\Delta)/P_{K,1}(\Delta) \simeq Gal(K_1/K).$$

From Proposition 2.5 it is clear that G_p splits completely in H_4 and H_4 contains $\sqrt{2}$. For any prime ideal π of K, $\left(\frac{K_1/K}{\pi}\right) = 1$. Hence, $\pi = x+y\sqrt{-7}$ is identity in $P_K(\Delta)/P_{K,1}(\Delta)$. Thus,

$$x + y\sqrt{-7} \equiv \pm 1 \bmod \Delta. \tag{1}$$

From Lemma 2.4, we have $4|y$. We assume that, $p > 7$, so $G_p \equiv 1 \pmod{32}$. Since, $4|y$ and $x \equiv \pm 1 \pmod 8$, we have,

$$x^2 + 7y^2 \equiv 1 \pmod{16}$$

and

$$(y \cdot \sqrt{-7})^2 \equiv 0 \pmod{16}.$$

That is $y \cdot \sqrt{-7} \equiv 0$ or $4 \pmod 8$. If $y \cdot \sqrt{-7} \equiv 4 \bmod 8$, then there is a contradiction to $\pi \equiv x + y\sqrt{-7} \equiv \pm 1 \bmod \Delta$ in (1) and to the splitting of π in H_4. Hence,

$$y \cdot \sqrt{-7} \equiv 0 \bmod 8 \tag{2}$$

Now, for $K = \mathbb{Q}(\sqrt{-7})$, let $\mathcal{O}_K$ be its ring of integers. The element $\omega = \frac{1+\sqrt{-7}}{2}$ is an integer of K, which is a zero of the polynomial $X^2 - X + 2$. As $-2 \in \mathbb{Z}/8\mathbb{Z}$ satisfies $X^2 - X + 2 = 0$, we obtain a ring homomorphism

$$\mathcal{O}_K \to \mathbb{Z}/8\mathbb{Z}$$

$$a + b\omega \to (a - 2b) \bmod 8.$$

Now $\sqrt{-7} = 2\omega - 1$ maps to 3, confirming $y \equiv 0 \pmod 8$ from equation 2. This completes the proof. □

2.2. **Artin's reciprocity law.** Now we give an alternate proof using the similar techinique used in [10] to prove Theorem 2.6.

2.3. **Primes in a Quadratic field.** In this section we refer mainly to [10]. To prove the results in the previous section, we need to consider the field $\mathbb{Q}(\sqrt{-7})$. We know that, for $K = \mathbb{Q}(\sqrt{-7})$, we obtain a ring homomorphism

$$\mathcal{O}_K \to \mathbb{Z}/8\mathbb{Z}$$

$$a + b\omega \to (a - 2b) \bmod 8.$$

kernel $\mathfrak{a}$ of the map is generated by 8 and $\omega + 2$. Thus $\mathfrak{a} = (\omega + 2)\mathcal{O}_K$. Their kernels are prime ideals of index 2 in $\mathcal{O}_K$ with generators ω and $\bar{\omega}$. Also, the identity $\omega\bar{\omega} = 2$ and $\omega + 2 = -\omega^3$ show that the ideal $\mathfrak{a}$ factors as the cube of prime $\omega\mathcal{O}_K$. Now consider an Abelian extension L of K with Galois group G, say $L = K[\beta]$ where β is a zero of $x^2 - \omega x - 1$ and L has dimension 2 over K. with $\beta^2 - \omega\beta - 1 = 0$. Now consider

$$K = \mathbb{Q}[\sqrt{-7}] = \mathbb{Q}[\omega] \text{ with } \omega^2 - \omega + 2 = 0 \text{ and}$$

$$L = K[\beta], \qquad \beta^2 - \omega\beta - 1$$

The discriminant of L/K is $\omega^2 + 4 = \omega + 2$ is non-zero and L has dimension 2 over K, hence L is Abelian over K with Galois group G of order 2, say $G = \{1, \rho\}$. The non-identity element ρ of G satisfies $\rho(\beta) = \omega - \beta = -1/\beta$.

The discriminant $\omega + 2 = -\omega^3$ of the polynomial defining L is not a square in K, hence we have $L = K[\sqrt{-\omega}]$.

The inclusion map $\mathcal{O}_K \to \mathcal{O}_L$ induces a ring homomorphism $k(\mathfrak{p}) \to \mathcal{O}_L/\mathfrak{p}\mathcal{O}_L$, where $k(\mathfrak{p}) = \mathcal{O}_K/\mathfrak{p}$, for a prime $\mathfrak{p}$ of K.

2.4. **An outline of the proof via Artin Symbol.** If $\mathfrak{p} = \pi\mathcal{O}_K$ is a prime of $K = \mathbb{Q}[\sqrt{-7}] = \mathbb{Q}[\omega]$ different from $\omega\mathcal{O}_K$, then the Artin symbol $\left(\frac{L/K}{\mathfrak{p}}\right) = 1$ or ρ according as π maps to ± 1 or to ± 3 under the map $\mathcal{O}_K \to \mathbb{Z}/8\mathbb{Z}$ that sends ω to -2. For example, $\sqrt{-7} = 2\omega - 1$ maps to 3 mod 8 and the number $(8k \pm 1) \pm 8l\sqrt{-7}$ maps to $\pm 1 \pmod 8$ for $l, k \in \mathbb{Z}$. This is because, $\sqrt{-7} = 2\omega - 1$ and ω maps to -2 under $\mathbb{Z}/8\mathbb{Z}$. We see that, for $p = 47$, $G_{47} = 140737471578113 = 5732351^2 + 7 \cdot 3925696^2$ is nothing but,

$$5732351 \pm 3925696\sqrt{-7} \equiv -1 \pm 3(0) \equiv -1 \bmod 8.$$

Consider an element of the form $x + \sqrt{-7}y$ of norm $x^2 + 7y^2$ which is a rational prime. The Artin symbol equals 1 if $x + 3y$ is $\pm 1 \bmod 8$, and ρ otherwise. From Lemma 2.4, it is clear that x leaves the remainder $\pm 1 \bmod 8$ and y is divisible by 4. Hence the Artin symbol is 1 if and only if y is divisible by 8. This observation is equivalent to the assertion that, *any prime of $K = \mathbb{Q}(\sqrt{-7})$ of norm G_p has trivial Artin symbol in the quadratic extension $L = K[\omega]$*. The proof works in the extension $N = K(\sqrt{-\omega}, \sqrt{-\bar{\omega}})$, which is the union of two conjugate quadratic extensions $L = K(\sqrt{-\omega})$ and $L = K(\sqrt{-\bar{\omega}})$. Further details can be filled in by following Theorem 2.6 and the last theorem in [10] and [14].

3. Generalization of Theorem 2.6

In this section we generalize the result of Theorem 2.6 for $K = \mathbb{Q}(\sqrt{-d})$ that has the same properties as $K = \mathbb{Q}(\sqrt{-7})$. In this case there exists a cyclic extension H_4 of $S = \mathbb{Q}(\sqrt{-2 \cdot d})$ and $\sqrt{2} \in H_4$, so that G_p splits completely in H_4. For this one requires $\left(\frac{2}{d}\right) = 1$ and $\left(\frac{-d}{G_p}\right) = 1$. Thus, with these conditions we state the general form of Proposition 2.5.

Proposition 3.1. *Let $d \equiv 3 \pmod 4$ be a square free integer and $d > 0$. Let G_p be a Gaussian Mersenne prime and $G_p = x^2 + dy^2$ for $d \equiv 3 \pmod 4$ exists and $\left(\frac{2}{d}\right) = 1 = \left(\frac{-d}{G_p}\right)$. Suppose that there exists a cyclic extension H_4 of $S = \mathbb{Q}(\sqrt{-2 \cdot d})$ with $[H_4 : S] = 4$, $H_4 \subset H(S)$ and $\sqrt{2} \in H_4$. Then G_p splits completely in H_4.*

Proof. Proof of this proposition is similar to Proposition 2.5. □

Thus, the generalized form of Theorem 2.6 is stated below, proof of this theorem is similar to the proof of Theorem 2.6.

Theorem 3.2. *Let $d \equiv 7 \pmod{24}$ be a square free integer and $d > 0$. Suppose G_p is a Gaussian Mersenne prime which splits completely in a cyclic extension H_4 of $S = \mathbb{Q}(\sqrt{-2 \cdot d})$. If $G_p = x^2 + dy^2$ for $d \equiv 7 \pmod{24}$ exists then $8|y$.*

4. Conclusion

In the case of Mersenne primes $2^p - 1$, whenever $2^p - 1$ is a quadratic residue of 7, then $2^p - 1$ takes the form $x^2 + 7y^2$. Franz Lemmermeyer made an observation that, in this representation x is divisible by 8 and y leaves the remainder ± 3 when divided by 8. This was later proved by H. W. Lenstra and P. Stevenhagen using Artin reciprocity law and presented this result on the occasion of birth centenary of Emil Artin on 3 March, 1998 at Universiteit van Amsterdam[10]. In [14], authors made similar observations for rational primes $N(\frac{\alpha^p-1}{\alpha-1})$, for $\alpha = 2 + \sqrt{2}$ and proved using Artin's reciprocity law. In this paper, we have made an observation that, if a Gaussian Mersenne prime is a quadratic residue of 7, then in the representation of G_p as $x^2 + 7y^2$, we have $x \equiv \pm 1 \pmod 8$ and y is divisible by 8 and proved this result using Artin reciprocity law and generalized this result for $d \equiv 7 \pmod{24}$.

5. Acknowledgement

The authors would like to thank Professor P. Stevenhagen, Universiteit Leiden, for his valuable insights of class field theory concepts. The authors would also like to thank Professor U.K.Anandavardhanan, IIT Bombay, for helping to improve earlier versions of this manuscript. The first author would like to acknowledge National Board for Higher Mathematics, Government of India for funding this work through Post Doctoral Fellowship.

References

[1] P.Berrizbeitia and B.Iskra, (2010) Gaussian Mersenne and Eisenstein Mersenne Primes, *Math. Comp.* **79**(271) (2010) 1779–1791.

[2] H.Cohen and P.Stevenhagen, Arithmetic of Number Rings in *Algorithmic Number Theory*, eds. Buhler, J. & Stevenhagen P., vol. 44 (MSRI, 2008), pp. 209–266.

[3] H.Cohen and P.Stevenhagen, Computational Class Field Theory, in *Algorithmic Number Theory*, eds. Buhler, J. & Stevenhagen P., vol. 44 (MSRI, 2008), pp. 497–534.

[4] H. Cohen, *A Course in Computational Algebraic Number Theory* (Springer-Verlag Berlin, 1996).

[5] H.Cohen, *Advanced Topics in Computational Number Theory* (Springer-Verlag New York, 2000).

[6] D.Cox, *Primes of the form $x^2 + ny^2$ Fermat, Class Field Theory and Complex Multiplication* (John Wiley & Sons, 1989).

[7] W.Jagy, (http://math.stackexchange.com/users/10400), *Cornacchia's Algorithm,* http://math.stackexchange.com/questions/577631 (version: 2013-11-22).

[8] W. Jagy, (http://math.stackexchange.com/users/10400), *Integers of the form $x^2 - ny^2$*, http://math.stackexchange.com/questions/530037

[9] F.Lemmermeyer, *Class Field Towers,* [Online] (2010), Available from (http://www.rzuser.uni-heidelberg.de)

[10] H.W.Lenstra and P.Stevenhagen, *Mersenne Primes and Artin Reciprocity*, *Nieuw Arch. Wisk.* (2000) **5**(1), pp. 44–54.

[11] B.Jansen, *Mersenne Primes and Class Field Theory.* Ph.D thesis, (Universiteit Leiden, Netherlands 2012).

[12] B.Jansen, *Mersenne Primes of the Form $x^2 + dy^2$*. Master's thesis, (Universiteit Leiden, Netherlands 2002).

[13] J.S.Milne, *Class Field Theory*, [Online] (2013) Available from http://www.jmilne.org/math.

[14] Sushma Palimar and B.R.Shankar, *Mersenne Primes in Real Quadratic Fields,* J. Integer Seq., (2012) **15** Article 12.5.6.
[15] J. Wordsworth, (math.stackexchange.com/users/32441). *Cornacchia's Algorithm*, http://math.stackexchange.com/questions/575862 (version: 2013-11-23).

E-mail address: sushmapalimar@gmail.com, sushma@csa.iisc.ernet.in, ad@csa.iisc.ernet.in

Dept. of Computer Science & Automation, Indian Institute of Science, Bangalore - 560012